


General Relativity (GR)

M1 - Physique 2023-2024



AE+GR (1907-1917)

*Paul-Antoine Hervieux
Unistra/IPCMS
hervieux@unistra.fr*

$g_{\mu\nu}$

$$S[g] = \frac{1}{16\pi G} \int \sqrt{-g} (R - 2\lambda) d^4x$$

///) Theory: $[g_{ij}]$ link between geometry and gravity

$$R_{ab} - \frac{1}{2} R g_{ab} + \lambda g_{ab} = 8\pi G T_{ab}$$

$$\ddot{x}^d + \Gamma_{ab}^d \dot{x}^a \dot{x}^b = 0$$

GR Field equations

“Geometry tells matter how to move, and matter tells geometry how to curve”

To find the equations determining a gravitational field, one must first determine the action S_g for this field. The equations are then obtained by varying the sum of the field and particle actions with respect to g_{ij} .

Field equations

- The S_g action, like the electromagnetic action S_{field} , must be expressed as a scalar integral $\int H \sqrt{-g} d\Omega$ extended over all space and between two values of the time component x^0 . We remind you that $d\Omega = dx^0 dx^1 dx^2 dx^3 = c dt dV$ and $g = -1$ for SR. $g \equiv [g_{ij}]$
- The gravitational field equations must not contain derivatives of field potentials of order greater than 2, as is the case for the electromagnetic field equations (A_α plays the role of g_{ij}): $\partial_\alpha F^{\alpha\beta} = \mu_0 J^\beta$ with $F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$.
- Since the field equations are obtained by varying the action, the H expression must not contain derivatives of g_{ik} greater than one. So H must contain only the tensor \mathbb{G} and the symbols Γ .
- Unfortunately, the quantities g_{ik} and Γ_{kl}^i alone cannot be used to form a scalar. ||

GR Field equations

- There is, however, a scalar R - the curvature of 4-space - which admittedly contains the metric tensor \mathbb{G} and its first and second derivatives, but only linearly.

It is possible to simplify the expression of S_g by eliminating the second derivatives $\partial^2 g_{ik} / \partial x^p \partial x^q$ involved in the R invariant. (*)

After this simplification, the integrand of S_g will contain only the tensor \mathbb{G} and the Christoffel symbols Γ . We have

$$\delta \int R \sqrt{-g} d\Omega = \delta \int H \sqrt{-g} d\Omega , \quad (1)$$

with

$$H(g, \partial g) = g^{ik} (\Gamma_{il}^m \Gamma_{km}^l - \Gamma_{ik}^l \Gamma_{lm}^m) . \quad (2)$$

- (*) The demonstration is quite difficult. It can be found in "Modern Geometry - Methods and Applications", Part I. The Geometry of Surfaces, Transformation Groups, and Fields (B. A. Dubrovin , A. T. Fomenko , S. P. Novikov). The main tool is the Stokes' theorem.

GR Field equations

We can therefore write:

$$\delta S_g = -\frac{c^3}{16\pi G} \delta \int H \sqrt{-g} d\Omega = -\frac{c^3}{16\pi G} \delta \int R \sqrt{-g} d\Omega, \quad (3)$$

where G is called the universal gravitational constant measured in $\text{N}\cdot\text{m}^2/\text{kg}^2$. Verify that S_g has the dimensions of an action.

Starting from $\delta S_g = -\frac{c^3}{16\pi G} \delta \int (H \sqrt{-g}) d\Omega$ we can easily obtain the Euler-Lagrange equations as

$$\delta S_g = -\frac{c^3}{16\pi G} \int \left\{ \frac{\partial(H\sqrt{-g})}{\partial g^{ik}} - \frac{\partial}{\partial x^l} \frac{\partial(H\sqrt{-g})}{\partial \frac{\partial g^{ik}}{\partial x^l}} \right\} \delta g^{ik} d\Omega. \quad (4)$$

After calculation, one finds:

$$\frac{1}{\sqrt{-g}} \left\{ \frac{\partial(H\sqrt{-g})}{\partial g^{ik}} - \frac{\partial}{\partial x^l} \frac{\partial(H\sqrt{-g})}{\partial \frac{\partial g^{ik}}{\partial x^l}} \right\} = \boxed{R_{ik} - \frac{1}{2} g_{ik} R}. \quad (5)$$

GR Field equations

Action of the matter

$$\delta S_m = \frac{1}{2c} \int T_{ik} \delta g^{ik} \sqrt{-g} d\Omega , \quad (6)$$

where T_{ik} is the stress-energy tensor of matter. Gravitational interaction only comes into play for bodies with sufficiently large masses. The expression $T_{ik} = (p + \epsilon)\bar{u}_i\bar{u}_k - pg_{ik}$ should be used, where ϵ is the energy density of the body (ϵ/c^2 is the mass density, i.e. the mass of the body's /"proper" unit volume, i.e. the volume in the frame of reference where the body element is at rest) and p is the pressure

Note: in the presence of electromagnetic fields T_{ik} is the sum of the total stress-energy tensor including the one coming from the electromagnetic field.

GR Field equations

$$\boxed{S_m = \frac{1}{c} \int \Lambda \sqrt{-g} d\Omega} \rightarrow \delta S_m = \frac{1}{c} \int \left\{ \frac{\partial(\Lambda \sqrt{-g})}{\partial g^{ik}} - \frac{\partial}{\partial x^l} \frac{\partial(\Lambda \sqrt{-g})}{\partial \frac{\partial g^{ik}}{\partial x^l}} \right\} \delta g^{ik} d\Omega . \quad (7)$$

If we define

$$\boxed{\frac{1}{2} \sqrt{-g} T_{ik} = \frac{\partial(\Lambda \sqrt{-g})}{\partial g^{ik}} - \frac{\partial}{\partial x^l} \frac{\partial(\Lambda \sqrt{-g})}{\partial \frac{\partial g^{ik}}{\partial x^l}} ,} \quad (8)$$

one get's

$$\delta S_m = \frac{1}{2c} \int \sqrt{-g} T_{ik} \delta g^{ik} d\Omega . \quad (9)$$

We have used that $\boxed{T_{i;k}^k = 0}$: the divergence of the stress-energy tensor is zero.

Einstein tensor

We start from the Bianchi identity ($\nabla_t R_{irs}^l \equiv R_{irs;t}^l$)

$$\boxed{\nabla_t R_{irs}^l + \nabla_s R_{itr}^l + \nabla_r R_{ist}^l = 0 .} \quad (10)$$

According to their tensor form, this second Bianchi identity is valid in any coordinate system and at any point of the Riemann space.

GR Field equations

Performing a first contraction on the Bianchi identity for $t = l$, we obtain

$$\nabla_l R_{irs}^l + \nabla_s R_{ilr}^l + \nabla_r R_{isl}^l = 0 . \quad (11)$$

Taking into account the definition $R_{ik} = \frac{\partial \Gamma_{ik}^l}{\partial x^l} - \frac{\partial \Gamma_{il}^k}{\partial x^k} + \Gamma_{ik}^l \Gamma_{lm}^m - \Gamma_{il}^m \Gamma_{mk}^k$ of the Ricci tensor and the equality $R_{rsl}^l = -R_{rls}^l$ one obtains

$$\nabla_l R_{irs}^l + \nabla_s R_{ir} - \nabla_r R_{is} = 0 . \quad (12)$$

Since the change in variance by means of g_{ik} is permutable with the covariant we have:

$$\nabla_s R_r^k = \nabla_s g^{ik} R_{ir} = g^{ik} \nabla_s R_{ir} . \quad (13)$$

Multiply relation (12) by g^{ik} and use the permutation property (13), we obtain

$$\nabla_l R_{rs}^{kl} + \nabla_s R_r^k - \nabla_r R_s^k = 0 . \quad (14)$$

Performing a second contraction with respect to indices k and s , we get:

$$\nabla_l R_{rk}^{kl} + \nabla_k R_r^k - \nabla_r R_k^k = 0 . \quad (15)$$

GR Field equations

After contraction, let's change the summation index l to index k in the first term of the previous equation. Moreover, with the equality $R_{rk}^{kl} = R_{kr}^{lk}$ we obtain

$$2\nabla_k R_r^k - \nabla_r R = 0 . \quad (16)$$

The latter expression can also be written as:

$$\nabla_k \left(R_r^k - \frac{1}{2} \delta_r^k R \right) = \nabla_k S_r^k = 0 . \quad (17)$$

The expression in brackets in the previous expression is a tensor, denoted S_r^k , whose covariant components are given by:

$$S_{ij} = g_{ik} S_j^k = g_{ik} \left(R_j^k - \frac{1}{2} \delta_j^k R \right) = R_{ij} - \frac{1}{2} g_{ij} R . \quad (18)$$

The S_{ij} tensor is called the Einstein tensor. Due to the symmetry of the Ricci tensor, the Einstein tensor is also symmetrical. According to (17), it verifies the identities:

$$\nabla_k S_r^k = 0 . \quad (19)$$

A tensor that satisfies relations of the form (19) identically is called a conservative tensor.

GR Field equations

Elie Cartan showed that if both of the following conditions apply:

i) the quantities S_{ij} depend only on the gravitational potentials g_{ij} and their first- and second-order derivatives, and are linear with respect to the second-order derivatives; ii) the tensor S_{ij} satisfies the conservation equations $\nabla_k S_r^k = 0$, then the only tensors satisfying both above conditions are given by the formula

$$S_{ij} = h \left[R_{ij} - \frac{1}{2} g_{ij} (R + k) \right], \quad (20)$$

where h and k are two constants.

Elie Cartan



(1869-1951)

Vacuum equations

Vacuum equations

In GR, a region where $T_{ik} = 0$, i.e. no matter, is called a "vacuum". Einstein's equations are

$$R_{ij} - \frac{1}{2}Rg_{ij} = 0 . \quad (21)$$

By contracting with g^{ij} we obtain $R = 0$ ($g_{ij}g^{ij} = \delta_i^i = 4$). Reintegrating into (21) gives:

$$R_{ij} = 0 . \quad (22)$$

In other words, in the absence of matter, the Ricci tensor is zero. This is not a sufficient condition for flatness, however, so in general space-time can be curved even in the absence of matter. Matter is therefore not enough to determine the gravitational field, just as charges are not enough to determine the electromagnetic field. A space-time in which the Ricci tensor is zero everywhere is called Einstein space-time. Therefore:

$$\text{Riemann tensor} = 0 \leftrightarrow \text{flat space} ; \text{ Ricci tensor} = 0 \leftrightarrow \text{empty space} . \quad (23)$$

These equations define general relativity. They are sufficient to describe gravitational waves, black holes, the expansion of the universe and the Big Bang, to found GPS technology...

Einstein equations with matter

Einstein equations with matter

so from the principle of least action we deduce $\delta S_m + \delta S_g = 0$ leading to

$$-\frac{c^3}{16\pi G} \int \left(R_{ik} - \frac{1}{2} g_{ik} R - \frac{8\pi G}{c^4} T_{ik} \right) \delta g^{ik} \sqrt{-g} d\Omega = 0, \quad (24)$$

hence, given that δg^{ik} are arbitrary one obtains

$$R_{ik} - \frac{1}{2} g_{ik} R = \frac{8\pi G}{c^4} T_{ik}, \quad (25)$$

or

$$R_i^k - \frac{1}{2} g_i^k R = \frac{8\pi G}{c^4} T_i^k. \quad (26)$$

These are the gravitational field equations or Einstein equations. We have used that $g_{il} g^{lk} = \delta_i^k$. g^{ik} is the inverse of g_{ik} .

Cosmological constant

Contacting on the indices i and k we find ($\delta_i^i = 4$, $R \equiv R_i^i$, $T \equiv T_i^i$)

$$R = -\frac{8\pi G}{c^4} T_i^k . \quad (27)$$

Consequently, the field equations can be recopied as

$$R_{ik} = \frac{8\pi G}{c^4} \left(T_{ik} - \frac{1}{2} g_{ik} T \right) . \quad (28)$$

Note: The equations of the gravitational field are not linear. As a result, the principle of superposition is not true for gravitational fields, as it is for the electromagnetic field in SR. However, it should be borne in mind that we're generally dealing with weak gravitational fields whose equations are linear to a first approximation. So the superposition principle remains legitimate with the same approximation.

Note: From (20) Einstein's equation can be rewritten as

$$\nabla_k S_r^k = 0 \leftrightarrow \nabla_k T_r^k = 0$$

$$R_{ik} - \frac{1}{2} R g_{ik} + \Lambda g_{ik} = \frac{8\pi G}{c^4} T_{ik} . \quad (29)$$

Λ is called the cosmological constant.

RG with differential forms

$$S_g = \frac{1}{2\chi} \int R^{ab} \wedge \star(V_a \wedge V_b).$$

Curvature 2-form:

$$R^{ab} = \frac{1}{2} R_{\mu\nu}^{ab} dx^\mu \wedge dx^\nu$$

Proof:

$$\begin{aligned} R^{ab} \wedge \star(V_a \wedge V_b) &= \frac{1}{2} R_{\mu\nu}^{ab} \frac{1}{2} V_a^\alpha V_b^\beta \eta_{\alpha\beta\rho\sigma} dx^\mu \wedge dx^\nu \wedge dx^\rho \wedge dx^\sigma \\ &= \frac{1}{4} R_{\mu\nu}^{ab} V_a^\alpha V_b^\beta \eta_{\alpha\beta\rho\sigma} \eta^{\mu\nu\rho\sigma} d^4x \sqrt{-g} \\ &= -\frac{1}{2} R_{\mu\nu}^{ab} V_a^\alpha V_b^\beta (\delta_\alpha^\mu \delta_\beta^\nu - \delta_\alpha^\nu \delta_\beta^\mu) d^4x \sqrt{-g} \\ &= -R d^4x \sqrt{-g} \end{aligned}$$

Stress-energy tensor

Energy-momentum tensor of a set of independent particles (no interaction between them) of masses m_a .

We define the mass density as

$$\mu = \sum_a m_a \delta(\vec{r} - \vec{r}_a), \quad (35)$$

where \vec{r}_a are the radius vectors of the particles. We have:

$$T^{ik} = \mu c \bar{u}^i \bar{u}^k \frac{ds}{dt} = \mu c^2 \sqrt{1 - \beta^2} \bar{u}^i \bar{u}^k. \quad (36)$$

Conservation law

We verify that

$$\partial_k \left(T^{(field)}_i{}^k + T^{(particles)}_i{}^k \right) = 0. \quad (37)$$

The meaning of this relation is that the sum of the energies and momenta of the field-particles are conserved.

Stress-energy tensor

Energy-momentum tensor for macroscopic bodies.

In addition to the energy-momentum (or stress-energy) tensor for a system of point particles we shall also need the expression for this tensor for macroscopic bodies which are treated as being continuous. We have

$$T^{ik} = (p + \epsilon)\bar{u}^i\bar{u}^k - pg^{ik} , \quad (38)$$

or

$$T^{ik} = (p + \epsilon)\bar{u}^i\bar{u}^k - p\delta^{ik} . \quad (39)$$

In the above expression $\epsilon \equiv \rho c^2$ is the energy density, p the pressure and $\bar{u}^i = \frac{\gamma}{c} \frac{dx^i}{dt}$ the dimensionless velocity.

Conservation law

$$T_{fluid}^{ik} = \left(\rho + \frac{p}{c^2}\right)u^i u^k - p\delta^{ik} . \quad (40)$$

When $v^2 \ll c^2$ one has $\nabla_\alpha T_{fluid}^{\alpha\beta} = 0 \rightarrow$

$$\left\{ \begin{array}{l} \frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot (\rho \vec{v}) = 0 \end{array} \right. \quad (41)$$

$$\left\{ \begin{array}{l} \frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla})\vec{v} = \frac{1}{\rho} \vec{\nabla} p + \vec{F}_{ext} . \end{array} \right. \quad (42)$$

The first equation is the continuity equation and is related to the mass conservation. The other three are linked to the Euler equation of fluid mechanics (conservation of momentum).

Einstein's approach

- Generalization of the Poisson and Laplace equations.

$$\Delta\phi = 4\pi G\rho \quad (o)$$

ϕ is the gravitational field produced by the mass density ρ

- The equations must be expressed as relations between tensors in space-time.

$$(o) \Leftrightarrow S_{\lambda\mu} = \chi T_{\lambda\mu} \quad (oo) \quad \chi : \text{constant}$$

- $T_{\lambda\mu}$ is purely mechanical and describes the state of energy and matter distribution at each point. It generalizes the second member of Poisson's equation.
- $S_{\lambda\mu}$ generalizes the left member of the Poisson equation. It is a purely geometrical quantity. By analogy with $\Delta\phi$ it contains gravitational potentials and their first- and second-order derivatives.
- We know that $\nabla_{\mu}T^{\mu\nu} = 0$. This is the law of conservation of energy and momentum.

$$(oo) \Rightarrow \nabla_{\mu}S^{\mu\nu} = 0$$